

# Colorado Allergy and Anaphylaxis Emergency Care Plan and Medication Orders

Student's Name: \_\_\_\_\_ D.O.B. \_\_\_\_\_ Grade: \_\_\_\_\_  
School: \_\_\_\_\_ Teacher: \_\_\_\_\_

Place child's  
photo here

**ALLERGY TO:** \_\_\_\_\_

**HISTORY:** \_\_\_\_\_

**Asthma:** ☐ YES (higher risk for severe reaction) – refer to their asthma care plan  
☐ NO

## ◇ STEP 1: TREATMENT ◇

### SEVERE SYMPTOMS: Any of the following:

**LUNG:** Short of breath, wheeze, repetitive cough  
**THROAT:** Tight, hoarse, trouble breathing/swallowing  
**MOUTH:** Swelling of the tongue and/or lips  
**HEART:** Pale, blue, faint, weak pulse, dizzy  
**SKIN:** Many hives over body, widespread redness  
**GUT:** Vomiting or diarrhea (if severe or combined with other symptoms)  
**OTHER:** Feeling something bad is about to happen, Confusion, agitation

### MILD SYMPTOMS ONLY:

**NOSE:** Itchy, runny nose, sneezing  
**SKIN:** A few hives, mild itch  
**GUT:** Mild nausea/discomfort

### 1. INJECT EPINEPHRINE IMMEDIATELY

#### 2. Call 911

- Ask for ambulance with epinephrine
  - Tell EMS when epinephrine was given
- #### 3. Stay with child and
- Call parent/guardian and school nurse
  - If symptoms don't improve or worsen give second dose of epi if available as instructed below
  - Monitor student; keep them lying down. If vomiting or difficulty breathing, put student on side

Give other medicine, if prescribed. (see below for orders) Do not use other medicine in place of epinephrine. **USE EPINEPHRINE**

#### 1. Stay with child and

- Alert parent and school nurse
- Give antihistamine (if prescribed)

2. If two or more mild symptoms present or symptoms progress **GIVE EPINEPHRINE** and follow directions in above box

**DOSAGE: Epinephrine:** inject intramuscularly using auto injector (check one): ☐ 0.3 mg ☐ 0.15 mg

☐ If symptoms do not improve \_\_\_\_ minutes or more, or symptoms return, 2<sup>nd</sup> dose of epinephrine should be given if available

**Antihistamine:** (brand and dose) \_\_\_\_\_

**Asthma Rescue Inhaler** (brand and dose) \_\_\_\_\_

Student has been instructed and is capable of carrying and self-administering own medication. ☐ Yes ☐ No

Provider (print) \_\_\_\_\_ Phone Number: \_\_\_\_\_

Provider's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

## ◇ STEP 2: EMERGENCY CALLS ◇

1. If epinephrine given, **call 911**. State that an anaphylactic reaction has been treated and additional epinephrine, oxygen, or other medications may be needed.

2. Parent: \_\_\_\_\_ Phone Number: \_\_\_\_\_

3. Emergency contacts: Name/Relationship \_\_\_\_\_ Phone Number(s) \_\_\_\_\_

a. \_\_\_\_\_ 1) \_\_\_\_\_ 2) \_\_\_\_\_

b. \_\_\_\_\_ 1) \_\_\_\_\_ 2) \_\_\_\_\_

### DO NOT HESITATE TO ADMINISTER EMERGENCY MEDICATIONS

I give permission for school personnel to share this information, follow this plan, administer medication and care for my child and, if necessary, contact our health care provider. I assume full responsibility for providing the school with prescribed medication and delivery/monitoring devices and release the school and personnel from any liability in compliance with their Board of Education policies.

Parent/Guardian's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

School Nurse: \_\_\_\_\_ Date: \_\_\_\_\_

Student Name: \_\_\_\_\_ DOB: \_\_\_\_\_

**Staff trained and delegated to administer emergency medications in this plan:**

1. \_\_\_\_\_

Room \_\_\_\_\_

2. \_\_\_\_\_

Room \_\_\_\_\_

3. \_\_\_\_\_

Room \_\_\_\_\_


Self-carry contract on file: ☐ Yes ☐ No

Expiration date of epinephrine auto injector: \_\_\_\_\_

Keep the child lying on their back. If the child vomits or has trouble breathing, place child on his/her side.

**AUVI-Q™ (EPINEPHRINE INJECTION, USP) DIRECTIONS**

1. Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.
2. Pull off red safety guard.
3. Place black end against mid-outer thigh.
4. Press firmly and hold for 5 seconds.
5. Remove from thigh.




**ADRENALICK® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR DIRECTIONS**

1. Remove the outer case.
2. Remove grey caps labeled "1" and "2".
3. Place red rounded tip against mid-outer thigh.
4. Press down hard until needle enters thigh.
5. Hold in place for 10 seconds. Remove from thigh.



**EPIPEN® AUTO-INJECTOR DIRECTIONS**

1. Remove the EpiPen Auto-Injector from the clear carrier tube.
2. Remove the blue safety release by pulling straight up without bending or twisting it.
3. Swing and firmly push orange tip against mid-outer thigh until it 'clicks'.
4. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove auto-injector from the thigh and massage the injection area for 10 seconds.



If this conditions warrents meal accomodations from food service, please complete the form for dietary disability if required by district policy.

Additional information: \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Adopted from the Allergy and Anaphylaxis Emergency Plan provided by the American Academy of Pediatrics, 2017